

Side effects of this medicine

Common side effects are reported as below. If any of them occurs, check with your doctor/pharmacist: somnolence, dry mouth, fatigue, constipation, weight gain, dizziness, headache, etc.

In rare cases, symptoms described below may be the sign of side effects indicated in brackets []. If they occur, stop taking your medicine and check with your doctor immediately.

- anxiety, agitation, fever [serotonin syndrome]
- fever, throat pain, muscular pain [agranulocytosis, neutropenia]
- convulsion [convulsion]
- general malaise, loss of appetite, yellowing in the skin or white of the eye [hepatic dysfunction, jaundice]
- convulsion, impaired consciousness, headache [syndrome of inappropriate secretion of antidiuretic hormone]
- fever, general malaise, erythema with edema in the center [muco-cutaneo-ocular syndrome, erythema multiform]

Other side effects not listed here may occur. Check with your doctor/pharmacist if you have any worrisome symptom.

Storage and other information

- Keep the medicine out of the reach of children. Store it away from direct sunlight, heat and moisture.
- Do not store the remainder and ask your pharmacist how to discard it.

For doctor use only: Day Month Year

For further information, ask your doctor/pharmacist. "Kanjamuke Iyakuhiin Guide" and Japanese package insert information (for medical professionals) is available on the website of Pharmaceuticals and Medical Devices Agency.